

arcalion[®] 200

Sulbutiamine

treatment for asthenia in all its forms

ARCALION has specific neurotropic action. This results in better physical resistance, improved intellectual performance and psychological balance. This new substance is related to thiamine with important structural changes which explain liposolubility and Arcalion's activity on specific cerebral structures.

Indications: Symptomatic treatment of functional asthenia, physical and intellectual overwork:

Memory disorders : difficulty in concentration :

Psychogenic impotence of recent onset :

Reactive asthenia :

Fatigue related to states of depression and anxiety :

Fatigue following infections and operations :

Fatigue in the post partum period :

Menopausal psychosomatic disorders :

Geriatric psychosomatic disorders :

Dosage: 2 tablets with breakfast - For adult use only.

The dose can be increased to 3 tablets daily in resistant cases

Contraindications : Hypersensitivity to sulbutiamine.

Precautions : Pregnancy : No fetal toxicity or malformation has been reported. However, follow-up of women exposed to Arcalion during pregnancy is insufficient to exclude such a risk. Therefore, as a precautionary measure, it is preferable not to use Arcalion during pregnancy. **Lactation :** in the absence of data concerning the diffusion into breast milk, treatment is not recommended during breastfeeding.

Side-effects : Occasionally : mild agitation in the elderly subjects, allergic skin reactions. The urine may sometimes give out an odour.

Overdosage : In case of overdosage, restlessness, euphoria, and tremor may appear.

These symptoms are transient and leave no sequelae.

ARCALION can safely be combined with other drugs.

Presentation

Box of 20 sugar-coated tablets.

Storage Conditions : Protect from direct light, heat & moisture.



Manufactured by:

Servier Research & Pharmaceuticals [Pakistan] (Pvt.) Ltd.

9-Km Sheikhpura Road, Lahore.

Under licence Les Laboratoires Servier - France